

Part III: Impurities in Luttinger liquids

1. Luttinger liquids

2. Impurity effects

S. Andergassen, T. Enss (Stuttgart)

3. Microscopic model

V. Meden, K. Schönhammer (Göttingen)

4. Flow equations

U. Schollwöck (Aachen)

5. Results

Phys. Rev. B **65**, 045318 (2002); Europhys. Lett. **64**, 769 (2003);

Phys. Rev. B **70**, 075102 (2004); Phys. Rev. B **71**, 041302 (2005);

Phys. Rev. B **71**, 155401 (2005); Phys. Rev. B **73**, 045125 (2006)


1. Luttinger liquids

One-dimensional interacting Fermi systems **without correlation gaps** are **Luttinger liquids**.

(1D counterpart of Fermi liquid in 2D or 3D)

One-dimensional electron systems:

- Complex chemical compounds containing chains
- Quantum wires (in heterostructures)
- Carbon nanotubes
- Edge states




(Dekker's group)

Electronic structure of 1D systems:


Dispersion relations:

$$\epsilon_k = k^2/2m \quad (\text{low carrier density})$$
$$\epsilon_k = -2t \cos k \quad (\text{tight binding})$$

"Fermi surface": 2 points $\pm k_F$



Dispersion relation near Fermi points:



approx. linear:

$$\xi_k = \epsilon_k - \epsilon_F = v_F (|k| - k_F)$$

Electron-electron interaction:

has stronger effects than in 2D and 3D systems:

no fermionic quasi-particles, Fermi liquid theory **not** valid.

Fermi liquid replaced by **Luttinger liquid**:

- only **bosonic** low-energy excitations
(collective charge/spin density oscillations)
- **power-laws** with non-universal exponents

⇒ **Luttinger liquid theory**

Review: T. Giamarchi: *Quantum physics in one dimension* (2004)

Bulk properties of Luttinger liquids:

- **Bosonic** low-energy excitations with **linear** dispersion relation


$$\xi_q^c = u_c q, \quad \xi_q^s = u_s q \quad (\text{charge and spin channel})$$

⇒ specific heat $c_V \propto T$

- **DOS** for single-electron excitations:

$$D(\epsilon) \propto |\epsilon - \epsilon_F|^\alpha$$

vanishes at Fermi level ($\alpha > 0$)



DOS in principle observable by photoemission or tunneling.

- Density-density correlation function $N(q)$:

finite for $q \rightarrow 0$ (compressibility)

divergent as $|q - 2k_F|^{-\alpha_{2k_F}}$ for $q \rightarrow 2k_F$

($\alpha_{2k_F} > 0$ for repulsive interactions)

\Rightarrow enhanced back-scattering ($2k_F$) from impurity.

For spin-rotation invariant (and spinless) systems all exponents can be expressed in terms of one parameter K_ρ .

Asymptotic low energy behavior (power-laws) of Luttinger liquids described by Luttinger model:

$$H_{\text{LM}} = \text{linear } \epsilon_k + \text{forward scattering interactions}$$

It is exactly solvable and scale-invariant (fixed point).

For spinless fermions only one coupling constant, parametrizing interaction between left- and right-movers:

$$H_I = g \int dx n_+(x) n_-(x)$$

2. Impurity effects

How does a **single non-magnetic impurity** (potential scatterer) affect properties of a Luttinger liquid?



Non-interacting system:

Impurity induces **Friedel oscillations** (density oscillations with wave vector $2k_F$)

DOS near impurity **finite** at Fermi level

Conductance reduced by a **finite** factor (transmission probability)

Kane, Fisher '92: impurity in **interacting** system (spinless Luttinger liquid)

- Weak impurity potential:

Backscattering amplitude V_{2k_F} generated by impurity grows as $\Lambda^{K_\rho-1}$ for decreasing energy scale Λ .

($K_\rho < 1$ for repulsive interactions; V_{2k_F} is "**relevant**" perturbation of pure LL)

\Rightarrow Low energy probes see **high barrier** even if (bare) impurity potential is weak!

- Weak link:



DOS at **boundary** of LL vanishes as $|\epsilon - \epsilon_F|^{\alpha_B} \Rightarrow$

Tunneling amplitude t_{wl} between two weakly coupled chains scales to zero as Λ^{α_B} with $\alpha_B = K_\rho^{-1} - 1 > 0$ at low energy scales.

(t_{wl} is "**irrelevant**" perturbation of split chain)

Hypothesis (Kane, Fisher):

Any impurity effectively "cuts the chain" at low energy scales and physical properties obey weak link or boundary scaling. \Rightarrow

DOS near impurity:

$$D_i(\epsilon) \propto |\epsilon - \epsilon_F|^{\alpha_B} \quad \text{for } \epsilon \rightarrow \epsilon_F \text{ at } T = 0$$

Conductance through impurity:

$$G(T) \propto T^{2\alpha_B} \quad \text{for } T \rightarrow 0$$

supported within effective bosonic field theory by:

refermionization (Kane, Fisher '92)

QMC (Moon et al. '93; Egger, Grabert '95)

Bethe ansatz (Fendley, Ludwig, Saleur '95)

3. Microscopic model

Spinless fermion model:



$$H_{\text{sf}} = -t \sum_j (c_{j+1}^\dagger c_j + c_j^\dagger c_{j+1}) + U \sum_j n_j n_{j+1}$$

Properties (without impurities):

- **exactly solvable** by Bethe ansatz
- **Luttinger liquid** except for $|U| > 2t$ at half-filling
- **charge density wave** for $U > 2t$ at half-filling

Impurity potential added to bulk hamiltonian H_{sf} :

general form:
$$H_{\text{imp}} = \sum_{j,j'} V_{j'j} c_{j'}^\dagger c_j$$

"**site** impurity":

$$H_{\text{imp}} = V n_{j_0} \quad (j_0 \text{ impurity site})$$

"**hopping** impurity":

$$H_{\text{imp}} = (t - t') (c_{j_0+1}^\dagger c_{j_0} + c_{j_0}^\dagger c_{j_0+1})$$

Later also **double barrier** (two site or hopping impurities)

4. Flow equations

Starting point (for approximations):

Exact hierarchy of differential flow equations for 1-particle irreducible vertex functions with infrared cutoff Λ :

$$\frac{d}{d\Lambda} \Sigma^\Lambda = \text{Diagram: a grey circle with two external lines and a loop on top labeled } S^\Lambda \text{ and } \Gamma^\Lambda \text{ below it.}$$

$$\frac{d}{d\Lambda} \Gamma^\Lambda = \text{Diagram: two grey circles connected by two arcs, top arc labeled } S^\Lambda \text{ and bottom arc labeled } G^\Lambda \text{, plus a grey circle with a loop on top labeled } \Gamma_3^\Lambda \text{ below it.}$$

etc. for $\Gamma_3^\Lambda, \Gamma_4^\Lambda, \dots$

where

$$G^\Lambda = [(G_0^\Lambda)^{-1} - \Sigma^\Lambda]^{-1}$$

$$S^\Lambda = [1 - G_0^\Lambda \Sigma^\Lambda]^{-1} \frac{dG_0^\Lambda}{d\Lambda} [1 - \Sigma^\Lambda G_0^\Lambda]^{-1}$$

Cutoff:

At $T = 0$ sharp **frequency** cutoff: $G_0^\Lambda = \Theta(|\omega| - \Lambda) G_0$

At **finite** T (discrete Matsubara frequencies) **soft** cutoff with width $2\pi T$

G_0 bare propagator without impurities and interaction

Approximations:

Scheme 1 (first order):

Approximate $\Gamma^\Lambda \approx \Gamma^{\Lambda^0}$ (ignore flow of 2-particle vertex)

$\Rightarrow \Sigma^\Lambda$ **tridiagonal** matrix in real space

$$\frac{d}{d\Lambda} \Sigma^\Lambda = \text{Diagram}$$

Flow equation very **simple**; at $T = 0$:

$$\frac{d}{d\Lambda} \Sigma_{j,j}^\Lambda = -\frac{U}{2\pi} \sum_{s=\pm 1} \sum_{\omega=\pm\Lambda} \tilde{G}_{j+s,j+s}^\Lambda(i\omega) \quad \frac{d}{d\Lambda} \Sigma_{j,j\pm 1}^\Lambda = \frac{U}{2\pi} \sum_{\omega=\pm\Lambda} \tilde{G}_{j,j\pm 1}^\Lambda(i\omega)$$

where $\tilde{G}^\Lambda(i\omega) = [G_0^{-1}(i\omega) - \Sigma^\Lambda]^{-1}$.

Kane/Fisher physics already qualitatively captured!

Scheme 2 (second order):

Neglect Γ_3^Λ ; approx. Γ^Λ by flowing **nearest neighbor** interaction U^Λ

\Rightarrow 1-loop flow for U^Λ ; flow of Σ^Λ as in scheme 1 with renormalized U^Λ

$$\frac{d}{d\Lambda} \Sigma_{j,j}^\Lambda = -\frac{U^\Lambda}{2\pi} \sum_{s=\pm 1} \sum_{\omega=\pm\Lambda} \tilde{G}_{j+s,j+s}^\Lambda(i\omega) \quad \frac{d}{d\Lambda} \Sigma_{j,j\pm 1}^\Lambda = \frac{U^\Lambda}{2\pi} \sum_{\omega=\pm\Lambda} \tilde{G}_{j,j\pm 1}^\Lambda(i\omega)$$

Works **quantitatively** even for rather big U

Derivation of flow equation (scheme 1):

Flow equation for **self-energy**:

$$\frac{d}{d\Lambda} \Sigma^\Lambda(1', 1) = -T \sum_{2, 2'} e^{i\omega_2 0^+} S^\Lambda(2, 2') \Gamma_0(1', 2'; 1, 2)$$

$$\frac{d}{d\Lambda} \Sigma^\Lambda = \text{Diagram}$$

Single-scale propagator

$$S^\Lambda = G^\Lambda [\partial_\Lambda (G_0^\Lambda)^{-1}] G^\Lambda = -\frac{1}{1 - G_0^\Lambda \Sigma^\Lambda} \frac{\partial G_0^\Lambda}{\partial \Lambda} \frac{1}{1 - \Sigma^\Lambda G_0^\Lambda}$$

Self-energy and propagator diagonal in frequency: $\omega_1 = \omega_{1'}$ and $\omega_2 = \omega_{2'}$.

Γ_0 frequency-independent \Rightarrow Σ **frequency-independent**.

Sharp frequency cutoff ($T = 0$): $G_0^\Lambda(i\omega) = \Theta(|\omega| - \Lambda) G_0(i\omega) \Rightarrow$

$$S^\Lambda(i\omega) = \frac{1}{1 - \Theta(|\omega| - \Lambda) G_0(i\omega) \Sigma^\Lambda} \delta(|\omega| - \Lambda) G_0(i\omega) \frac{1}{1 - \Theta(|\omega| - \Lambda) \Sigma^\Lambda G_0(i\omega)}$$

$\delta(\cdot)$ meets $\Theta(\cdot)$: ill defined!

Consider regularized (smeared) step functions Θ_ϵ with $\delta_\epsilon = \Theta'_\epsilon$,
then take limit $\epsilon \rightarrow 0$, using

$$\int dx \delta_\epsilon(x - \Lambda) f[x, \Theta_\epsilon(x - \Lambda)] \xrightarrow{\epsilon \rightarrow 0} \int_0^1 dt f(\Lambda, t) \quad \text{proof: substitution } t = \Theta_\epsilon$$

Integration can be done analytically, yielding

$$\frac{d}{d\Lambda} \Sigma_{j'_1, j_1}^\Lambda = -\frac{1}{2\pi} \sum_{\omega = \pm\Lambda} \sum_{j_2, j'_2} e^{i\omega 0^+} \tilde{G}_{j_2, j'_2}^\Lambda(i\omega) \Gamma_{j'_1, j'_2; j_1, j_2}^0$$

where $\tilde{G}^\Lambda(i\omega) = [G_0^{-1}(i\omega) - \Sigma^\Lambda]^{-1}$

Insert real space structure of **bare vertex** for spinless fermions with nearest neighbor interaction U :

$$\Gamma_{j'_1, j'_2; j_1, j_2}^0 = U_{j_1, j_2} (\delta_{j_1, j'_1} \delta_{j_2, j'_2} - \delta_{j_1, j'_2} \delta_{j_2, j'_1})$$

$$U_{j_1, j_2} = U (\delta_{j_1, j_2-1} + \delta_{j_1, j_2+1})$$

\Rightarrow Flow equations

$$\frac{d}{d\Lambda} \Sigma_{j,j}^\Lambda = -\frac{U}{2\pi} \sum_{s=\pm 1} \sum_{\omega=\pm\Lambda} e^{i\omega 0^+} \tilde{G}_{j+s, j+s}^\Lambda(i\omega)$$

$$\frac{d}{d\Lambda} \Sigma_{j, j\pm 1}^\Lambda = \frac{U}{2\pi} \sum_{\omega=\pm\Lambda} e^{i\omega 0^+} \tilde{G}_{j, j\pm 1}^\Lambda(i\omega)$$

Convergence factor $e^{i\omega 0^+}$ matters only for $\Lambda \rightarrow \infty$

Initial condition at $\Lambda = \Lambda_0 \rightarrow \infty$:

$$\Sigma_{j_1, j'_1}^{\Lambda_0} = V_{j_1, j'_1} + \frac{1}{2} \sum_{j_2} \Gamma_{j'_1, j_2; j_1, j_2}^0$$


where V_{j_1, j'_1} is the bare impurity potential and the second term is due to the flow from ∞ to Λ_0 (!)

Flow equations at finite temperatures $T > 0$:

Replace $\omega = \pm\Lambda$ by $\omega = \pm\omega_n^\Lambda$ in flow equations, where ω_n^Λ is the Matsubara frequency most close to Λ .

Calculation of **conductance**:

Interacting chain connected to semi-infinite **non-interacting leads** via smooth or abrupt **contacts**



Conductance $G(T) = -\frac{e^2}{h} \int d\epsilon f'(\epsilon) |t(\epsilon)|^2$ with $|t(\epsilon)|^2 \propto |G_{1,N}(\epsilon)|^2$

Propagator $G_{1,N}(\epsilon)$ calculated in presence of **leads**, which affect the interacting region only via **boundary contributions** $\Sigma_{1,1}(\epsilon)$ and $\Sigma_{N,N}(\epsilon)$ to the self-energy

Vertex corrections vanish within our approximation (no inelastic scattering)


(see Oguri '01)

fRG features:

- perturbative in U (weak coupling)
- **non-perturbative** in impurity strength
- **arbitrary** bare impurity potential (any shape)
- **full** effective impurity potential
(cf. **Matveev, Yue, Glazman '93**: only V_{2k_F})
- cheap numerics up to 10^5 sites for $T > 0$ and 10^7 sites at $T = 0$.
- captures **all scales**, not just asymptotics.

5. Results

Renormalized impurity potential (from self-energy Σ_{jj} at $\Lambda = 0$):




long-range $2k_F$ -oscillations ! (associated with Friedel oscillations of density)

$2k_F$ -oscillations also in renormalized hopping amplitude around impurity

Results for local DOS near impurity site:

(half-filling, ground state, $U = 1$, $V = 1.5$, 1000 sites)



Strong **suppression** of DOS near Fermi level

Power law with boundary exponent α_B for $\omega \rightarrow 0$, $N \rightarrow \infty$

Spectral weight at $\omega = 0$ in good agreement with DMRG for $U < 2$.

Log. derivative of **spectral weight**
at Fermi level as fct. of system size:

- near **boundary** (*solid lines*)
- near **hopping impurity** (*dashed lines*)

circles: quarter-filling, $U = 0.5$

squares: quarter-filling, $U = 1.5$

open symbols: **fRG**



filled symbols: **DMRG**

top panel: **without** vertex renorm.


bottom panel: **with** vertex renorm.

horizontal lines:

exact **boundary exponents**





Friedel oscillations from open boundaries:
(half-filling, ground state)




Excellent agreement between **fRG** and **DMRG**

One parameter scaling of conductance ($T = 0$):



Single impurity, smooth contacts: $G(N) = \frac{e^2}{h} \tilde{G}_K(x)$, $x = [N/N_0(U, V)]^{1-K}$



Crossover size
as function of bare
reflection amplitude





Conductance at $T > 0$ — smooth contacts



Asymptotic power law $G(T) \propto T^{2\alpha}$ reached on accessible scales only for sufficiently strong impurities


Resonant tunneling through double barrier:




(Dekker's group '01)

Treated theoretically by many groups; **controversial** results !

Model setup:





Resonance peaks in conductance as a function of gate voltage:



At $T = 0$, width $w \sim N^{K-1}$

T -dependence of $|t(\epsilon)|^2$ important

fRG results for $G_p(T)$ (symmetric double barrier):



Various distinctive **power laws**, in particular (Furusaki, Nagaosa '93,'98):

- exponent $2\alpha_B$ (looks like independent impurities in series)
- exponent $\alpha_B - 1$ ("uncorrelated sequential tunneling")

No indications of exponent $2\alpha_B - 1$ ("correlated sequential tunneling")

Summary . . .

- fRG is reliable and flexible tool to study Luttinger liquids with impurities
- can be applied to microscopic models, restricted to "weak" coupling
- provides simple physical picture
- interplay of contacts, impurities, and correlations
- method covers all energy scales
- resonant tunneling: universal behavior and crossover captured

. . . and outlook

- include **spin**
(extended Hubbard model: [Andergassen et al., PRB 73, 045125 \(2006\)](#))
- more complex geometries
(Y-junctions: [Barnabé-Thériault et al., PRL 94, 136405 \(2005\)](#))
- include bulk anomalous dimension
- include inelastic processes
- extend to non-linear transport